

CPO prediction: accuracy assessment and impact on UT1 intensive results

Zinovy Malkin

Pulkovo Observatory, St. Petersburg, Russia

Abstract

The UT1 Intensives results obtained from single-base observations heavily depend on the celestial pole offset (CPO) model used during data processing. Since accurate CPO values can be obtained only from the 24h VLBI sessions and are available with delay from two to four weeks, CPO predictions are necessarily applied to the UT1 intensive data analysis, and errors in predictions can influence the operational UT1 accuracy. In this paper the real accuracy of CPO prediction is assessed using the actual IERS and PUL predictions made in 2007-2009. Also, results of operational processing was analyzed to investigate the actual impact of EOP prediction errors on the rapid UT1 results. It was found that the impact of CPO prediction errors is at a level of several microseconds, whereas the impact of the inaccuracy in the polar motion prediction may be about one order larger for ultra-rapid UT1 results. The situation can be amended if the IERS Rapid solution will be updated more frequently.

1 Introduction

Rapid and ultra rapid VLBI UT1 observations are vital for accuracy of the rapid IERS EOP solution and its prediction. To decrease rapid UT1 latency, the special single-base 1-hour sessions are conducted practically every day with delay of processing from several hours to several days. As was shown in previous studies [1–3] UT1 estimates obtained from the single-base intensive programs heavily depend on the celestial pole motion model used during analysis. For the most exacting applications, the celestial pole coordinates are computed as the sum of the theoretical values given by a adopted theory of precession-nutations, IAU2000A nowadays, and corrections called celestial pole offset (CPO) and obtained from observations, exclusively VLBI nowadays. The CPO comprises of trends and (quasi)periodic components, Free Core Nutation (FCN) is the first place, caused by the inaccuracy of Earth Rotation theory.

The most accurate CPO can be obtained only from the 24h VLBI sessions and are available, as a rule, with delay from two to four weeks¹. Therefore CPO predictions are necessarily applied to the UT1 intensive data analysis, and errors in predictions can influence the rapid UT1 accuracy. In this papers the real accuracy of CPO prediction is assessed using the actual predictions made by IERS (USNO) and PUL IVS Analysis Center (Pulkovo observatory). The required prediction length can be found from analysis of the IVS combination delay, i.e. the time between the date of publication of IVS combined solution and the last EOP epoch in this solution (see Fig 1). Of course, IVS series is then updated with new observations processed, but this changes in the IVS data are small enough to significantly influence rapid UT1 results. One can see that the required length of CPO forecast is about 40 days. We extend our analysis to longer length, which may be interesting for other applications.




Figure 1: Delay of IVS combined EOP.

This paper is aimed at accuracy assessment of the CPO predictions computed with different models. As usually, the prediction accuracy is derived from comparison of predicted values with the final ones. For proper interpretation of the results obtained in this study, the following circumstance should be taken into account. Each CPO model is a result of fitting of observed CPO series. The models differ not only by method of fitting, but also by CPO data used for analysis, which makes results of accuracy assessment somewhat ambiguous. One may consider the prediction accuracy with respect to the model itself, which is, in fact, the accuracy of representation of given model, which can be called internal prediction accuracy of the model. However, we are interested in the accuracy of representation of the actual celestial pole motion, which is most

¹Strictly speaking, CPO results from individual analysis center are available with lower delay, but we consider the IVS combined CPO series as the most suitable for the EOP service applications

important for majority of users. For this reason we use a comparison of CPO prediction made by various authors with the IVS combined CPO series, which is intended to be an official IERS standard. The internal prediction errors were also computed for models used in this work, but do not discussed here.

2 Data used

In this study we present results of processing of VLBI observations made in the framework of the UT1 intensives IVS observing program with different delay and different CPO models. The following data were used:

- INT1 sessions, observed on the workdays on the stations KOKEE (Kk) and WETTZELL (Wz); database is normally available in 2–5 days.
- INT2 sessions, observed at weekends on the stations TSUKUB32 (Ts) and WETTZELL; database is normally available in 1–2 days.
- INT3 sessions, observed on Monday on the stations NYALES20 (Ny), TSUKUB32, and WETTZELL; database is normally available in the same day.

The following actual and publicly available CPO models were tested:

- IERS final EOP series computed at the Paris Observatory (C04 series) [4]. It does not contain prediction and thus is equivalent to zero model for rapid data processing.
- IERS rapid EOP series computed at the USNO (NEOS model) [4]. It is constructed from analysis of the NEOS combined CPO series and updated daily.
- Lambert’s FCN series computed at the Paris Observatory (SL model) [5]. As a matter of fact, it can represent only the FCN contribution to CPO. However, this model is recommended by the IERS Conventions (2003) as the substitute for CPO. It is constructed from analysis of the IERS combined series C04 and updated every several months.
- Author’s ZM2 model computed at the Pulkovo Observatory [6]. It is constructed from analysis of the IVS combined CPO series and updated daily.

Comparison of this models with IVS data is shown in Fig. 2.




Figure 2: CPO models compared with the IVS combined CPO series.

3 Accuracy of CPO predictions

As usually, the accuracy of CPO predictions was estimated from comparison of predicted and final values. Predictions made in the period from December 30, 2006 till December 25, 2009 were used. Both rms and maximum prediction error was computed; the latter caused the maximum dilution in the UT1 accuracy, and thus is very important. Results are presented in Figs 3 and 4. One can see that ZM2 model provides the best accuracy of CPO prediction.




Figure 3: The rms error of CPO prediction for different models.




Figure 4: The maximum absolute error of CPO prediction for different models.

4 Impact on rapid UT1

In this section we investigate the impact of EOP prediction on the rapid UT1 results. To investigate the impact of the CPO prediction error on the rapid UT1 estimate we started in October 2009 experimental processing of INT1, INT2, and INT3 sessions with four aforementioned CPO models. All the computations presented here were made with ZM2 model. Using other models gives similar results. This experiment started with the session I09295 observed on October 22, 2009. Each session is processed processed triply with different delay:

1. immediately after the database is available (O); during INT1 and INT2 processing interpolated polar motion (PM) and extrapolated CPO are used, during INT3 processing both CPO and PM are extrapolated;
2. 5-7 days after the date of observations (O2); during this processing practically final PM is available, but CPO is still extrapolated;
3. at least 10 days after the IVS combined CPO for the date is available (F); during this processing both final CPO and PM are available.

In the beginning of this work, only operational processing were performed. Correspondingly, the first final processing was made in December 2009. Comparison of the final and operational results showed large difference between them for some sessions, especially for INT3 that were operationally processed before the IERS PM solution is available for the date. To separate the impact of the errors in CPO and polar motion, starting with the session K09346 (December 12, 2009) the O2 estimates are computed.

The differences between the UT1 estimates obtained with different delays are shown in Tables 1 and 2. Comparison of F-O, O2-O, and F-O2 differences shows that errors in extrapolated PM coordinates has more significant impact on the UT1 estimates then CPO prediction errors. This can be explained by the fact that while the maximum CPO 30-day prediction error during the last three months was about 0.15 mas, the maximum PM error for the IERS Bulletin A 1-day prediction during the same period was about 1.7 mas and 0.94 mas for X and Y pole coordinates respectively. Consequently, the impact of the PM prediction errors is about one order larger than the impact of the CPO prediction errors. The latter is at a level of a few microarcseconds, much less than the uncertainty of UT1 estimates. This result is in good agreement with Nothnagel's estimate of single-base UT1 bias at the level of 1 microsecond for the 40-microarcsecond bias in CPO or PM [2].

One can see that INT3 results obtained for all three stations NyTsWz are similar to the single-base solutions TsWz for the same sessions.

Table 1: Differences between the UT1 estimates obtained with different delays: a case of the KkWz baseline. See explanation in text. Unit: microseconds

Session code	Week-day	UT1 differences			Session code	Week-day	UT1 differences		
		F-O	F-O2	O2-O			F-O	F-O2	O2-O
I09295	Thu	-0.8			I09348	Mon	-2.2	-2.2	0.0
I09296	Fri	-0.8			I09349	Tue	-2.1	-1.6	-0.5
I09299	Mon	-0.8			I09350	Wed	-0.2	-1.3	1.1
I09300	Tue	-1.6			I09351	Thu	-0.8	-0.8	0.0
I09302	Thu	-1.6			I09352	Fri	-0.9	-0.8	-0.1
I09303	Fri	-1.7			I09355	Mon	-1.3	-1.3	0.0
I09306	Mon	-0.3			I09356	Tue	-1.6	-1.6	0.0
I09308	Wed	-2.6			I09357	Wed	-2.0	-2.0	0.0
I09309	Thu	-1.8			I09362	Mon	-1.5	-1.6	0.1
I09310	Fri	-1.6			I09363	Tue	-1.7	-1.5	-0.2
I09313	Mon	-1.7			I09364	Wed	-1.5	-1.1	-0.4
I09314	Tue	-1.4			I09365	Thu	-1.9	-1.6	-0.3
I09315	Wed	-1.0			I10004	Mon	-0.3	-1.6	1.3
I09316	Thu	3.0			I10005	Tue	-1.8	-1.8	0.0
I09317	Fri	-2.0			I10006	Wed			-0.4
I09320	Mon	-2.5			I10007	Thu			-0.5
I09321	Tue	-2.5			I10008	Fri			0.1
I09322	Wed	-2.7			I10011	Mon			0.7
I09323	Thu	-2.8			I10012	Tue			-0.2
I09324	Fri	-2.9			I10013	Wed			0.1
I09327	Mon	-2.3			I10014	Thu			0.0
I09328	Tue	-2.4			I10015	Fri			-0.2
I09329	Wed	-2.7			I10019	Tue			-0.1
I09334	Mon	-4.1			I10021	Thu			-0.6
I09335	Tue	-2.8			I10022	Fri			-0.3
I09336	Wed	-2.2			I10025	Mon			0.8
I09337	Thu	-2.6			I10026	Tue			1.4
I09338	Fri	-3.1			I10027	Wed			2.0
I09341	Mon	-2.7							
I09342	Tue	-2.7							
I09344	Thu	-2.5							
I09345	Fri	-2.5							

Table 2: Differences between the UT1 estimates obtained with different delays: a case of the TsWz and NyTsWz baselines. See explanation in text. Unit: microseconds.

Session code	Week-day	UT1 differences					
		TsWz			NyTsWz		
		F-O	F-O2	O2-O	F-O	F-O2	O2-O
K09299	Mon	-1.6					
K09304	Sat	-3.4					
K09305	Sun	9.2					
K09306	Mon	46.9					
K09311	Sat	-3.4					
K09312	Sun	1.3					
K09313	Mon	8.1					
K09318	Sat	1.1					
K09319	Sun	-0.8					
K09320	Mon	-0.2					
K09325	Sat	-2.0					
K09326	Sun	-1.5					
K09327	Mon	20.2					
K09332	Sat	0.2					
K09333	Sun	5.3					
K09334	Mon	-2.0					
K09339	Sat	-1.2					
K09340	Sun	0.5					
K09341	Mon	-1.0					
K09346	Sat	-0.3	0.7	-1.0			
K09347	Sun	0.5	1.9	-1.4			
K09348	Mon	-9.0	0.3	-9.3			
K09353	Sat	-0.6	0.1	-0.7			
K09354	Sun	0.9	0.7	0.2			
K09355	Mon	4.1	0.0	4.1			
K09360	Sat	0.2	0.2	0.0			
K09361	Sun	2.8	-0.2	3.0			
K10002	Sat	-1.3	-1.2	-0.1			
K10003	Sun	0.6	-0.9	1.5			
K10004	Mon	4.4	-0.2	4.6	2.1	-0.5	2.6
K10009	Sat			-0.3			
K10010	Sun			-0.2			
K10011	Mon			8.7			9.0
K10016	Sat			2.4			
K10017	Sun			0.7			
K10018	Mon			-12.5			-10.6
K10023	Sat			0.5			
K10024	Sun			0.5			
K10025	Mon			-2.2			-2.1

5 Conclusion

The impact of CPO prediction error on the rapid UT1 results seems to be not very significant, much less than the impact of the PM prediction error. The most rapid UT1 observations of the INT3 observing program that are correlated in the day of observations, so that the database is normally available before the IERS Rapid combination used as a priori EOP is updated, show sometimes very large bias up to several tens microseconds as compared with result of processing made after interpolated IERS PM data is published. The situation can be amended if the IERS Rapid solution will be computed and published more frequently, say every 6 hours, after the ultra-rapid IGS combination is ready. Such an approach seems to be much more preferable to a user's home-bred combination of the IERS and IGS data.

References

- [1] Titov O.A. (2000) Influence of Adopted Nutation Model on VLBI NEOS-Intensives Data Analysis. In: Proc. IAU Coll. 180, Washington DC, 27–30 March 2000, Eds. K.J.Johnston, D.D.McCarthy, B.J.Luzum, G.H.Kaplan, 259–262.
- [2] Nothnagel A. and Schnell D.(2008) The impact of errors in polar motion and nutation on UT1 determinations from VLBI Intensive observations. *J. Geodesy*, V. 82, No. 12, 863–869.
- [3] Malkin Z. (2009) A possible improvement of the VLBI Intensives UT1 results using FCN models. Presented at the IERS Workshop on EOP Combination and Prediction, Warsaw, 19-21 Oct 2009, http://www.cbk.waw.pl/EOPPW2009/contributions/session1/session1.1/mon10_Malkin.pdf, visited on January 25, 2010.
- [4] IERS Annual Report 2007, Verlag des Bundesamts für Kartographie und Geodäsie, Frankfurt am Main, 2009.
- [5] Lambert S.B. (2009) Empirical Model of the Free Core Nutation (Technical Note). <http://syrte.obspm.fr/~lambert/fcn/notice.pdf>, visited on January 25, 2010.
- [6] Malkin Z.M. (2007) Empiric Models of the Earth's Free Core Nutation. *Solar System Research*, V. 41, No. 6, 492–497.